



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

'Ornithology of a Churchyard,' by B. S. Bowdish.

As the opening paper of the afternoon, Mr. Dutcher, Chairman of the 'Committee on Protection of North American Birds,' read a summary of the report of his committee for the previous year.

The concluding paper, illustrated by lantern slides, was :

'The Season's Experiences with Shore-birds, Herons and Water-fowl,' by Rev. H. K. Job.

The next meeting of the Union will be held in New York City, in November, 1905.

The good attendance at this Congress, together with the new members elected, the largest number in any year since the founding of the Union, was most gratifying.

JNO. H. SAGE,
Secretary.

GENERAL NOTES.

Sabine's Gull in Montana.—I enclose two photographs of an immature example of Sabine's Gull (*Xema sabinii*), which, as far as I know, has not before been recorded from Montana. This bird was shot September 23, on the Yellowstone, by the ferryman at Terry, who told me that on the previous day he had seen about fifty gulls, all resting on the river shore—the flock including this species and others slightly larger. I presumed the larger gulls to be Ring-billed Gulls, which, as well as American Herring Gulls, I have observed on the Yellowstone at different dates.

The specimen is a male. The dimensions in inches are: Length to end of tail, 13; to end of wings, 14; extent, 30; wing, 10; tail, 5; across fork of tail, 1½; bill, along gape, 1½. The bill was black; legs and feet flesh color; irides dark brown.—E. S. CAMERON, *Terry, Montana*.

Additional Record of the European Widgeon (*Marcca penelope*).—I have recently examined a fine adult male of this species, which was shot on an overflowed meadow near Nippersink Lake, Lake County, Ill., on April 1, 1904, by Mr. Charles Muehrcke, and is now in his possession. The bird was in company with six of his American cousins, all of whom were shot. The specimen is mounted representing dead game. This record makes the eighteenth for the interior.—RUTHVEN DEANE, *Chicago, Ill.*

Little Blue Heron in Connecticut.—A flock of Little Blue Herons (*Florida cærulea*), all young in the white plumage, made its headquarters